

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Ants, Nuisance	acephate	Orthene TT&O 75 and 97	1.2 to 1.6 oz/gal, see label	<p>In most cases ants serve as predators of turfgrass pests and competitors to red imported fire ants. Ants become a nuisance when they build up to a high number, invade buildings and equipments, and build mounds that interfere with the smoothness of the greens. Species identification and an understanding of ant biology are critical in deciding where to treat and what to treat with, especially when baits are used. Ants are very susceptible to insecticide treatments, but relief of the problem seldom occurs unless the colony itself is eliminated. Sap-sucking insects (such as aphids, mealybugs, scale insects and leafhoppers) on nearby vegetations or landscape ornamentals should be controlled to reduce their attraction to the honeydew-seeking ants.</p> <p>Check labels of Sevin products for site restrictions.</p> <p>Dursban PRO is for uses on golf courses, road medians, and industrial plants sites only. Dursban 50W can also be used on sod farms and seed productions.</p> <p>Use GC formulation of Aloft for golf course and sod farms. Use LC formulation of Aloft for residential and commercial lawns, parks, recreational areas, and athletic fields.</p> <p>Check label for site restrictions.</p> <p>Demon is for lawn and landscape uses.</p> <p>DeltaGard is for lawns, recreational areas and athletic fields. Use GC formulation for golf courses and sod farms.</p> <p>For fire ants and big-headed ants. Not for use on sod farms and seed productions.</p> <p>Broadcast or slit applications. Provide 3 months control. Not for pasture and grazing lands. Check label for buffer zone and yearly application limit requirements.</p> <p>For fire ants, big-headed ants, and harvester ants.</p> <p>Use GC formulation of Allectus on golf courses and sod farms.</p> <p>For fire ants, big-headed ants, and pavement ants.</p> <p>Demand and Scimitar for use on lawns, recreational areas and athletic fields. GC formulation also for gold course and sod farm uses.</p> <p>For fire ants, native ants, big-headed ants, harvester ants, and Argentine ants.</p> <p>Astro is for use on lawns, recreational areas and athletic fields.</p> <p>For fire ants and big-headed ants.</p>
	bifenthrin ²	Onyx	0.07 to 0.15 fl oz	
		OnyxPro	0.16 to 0.32 fl oz	
		Talstar EZ Golf , GC Gran, PL	2.3 to 4.6 lbs	
		Talstar GC Flowable, Talstar One	0.25 to 1.0 fl oz	
	carbaryl	Sevin 5 Bait	11 oz	
		Sevin 10G	1.4 to 1.9 lbs	
		Sevin 80 WSP	2.5 to 5.0 lb/acre	
		Sevin SL	1.5 to 3 fl oz	
	chlorpyrifos	Dursban 50W	2 lb/acre	
		Dursban PRO	1.5 fl oz	
	clothianidin	Arena .25G	1.84 to 3.67 lbs	
		Arena .5G	1.0 to 1.8 lbs	
	clothinidin + bifenthrin	Aloft GC SC, LC SC	11.65 to 23.3 fl oz, see label	
		Aloft GC G, LC G	80 to 160 lbs, see label	
	cyfluthrin	Tempo (various formulations)	See label	
	cypermethrin	Demon Max	0.5 fl oz/gal	
		Demon WP	0.33 oz/gal	
	deltamethrin	DeltaGard GC, DeltaGard T&O	0.4 to 0.6 fl oz	
		DeltaGard G, DeltaGard GC Gran	2 to 3 lbs	
	fenoxycarb (bait)	Award Fire Ant Bait	1 to 1.5 lbs/acre	
	fipronil	Chipco TopChoice	2 lbs	
		Chipco Choice	4.6 oz	
	hydramethylnon (bait)	Amdro Pro, SiegePro	1 to 1.5 lbs/acre	
		Maxforce G	1 to 2 lbs	
	imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	1.7 to 2.9 lbs	
	Allectus SC, Allectus GC SC	1.32 to 1.65 fl oz		
indoxacarb (bait)	Advion Fire Ant Bait	1.5 lb/acre		
lambda-cyhalothrin	Demand EZ	13.6 to 28 ml		
	Demand G	2 to 3 lbs		
	Demand CS, Scimitar GC and CS	3.4 to 7 ml		
methoprene + hydramethylnon (bait)	Extinguish Plus	1.5 lbs/acre		
permethrin	Astro	0.4 to 0.8 fl oz		
pyriproxyfen (bait)	Distance Fire Ant Bait	1 to 1.5 lbs/acre		

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments	
Ants, Red Imported Fire	Mound treatment and Broadcast insecticide				
	acephate	Orthene TT&O 75	Drench: 0.2 oz/gal/mound Dry: 1-2 teaspoons/mound	<p>The Red Imported Fire Ant (RIFA) found in South Carolina is an invasive species. Movement of soil and plant materials is regulated by a federal quarantine. If shipment of soil, sods or plant crops outside of the quarantine area is intended see http://www.aphis.usda.gov/oa/pubs/ifapub.pdf for regulations on the specific treatments required by USDA-APHIS, or consult with the SC Department of Plant Industry.</p> <p>Chemicals for RIFA management can be broken into three categories: Individual mound treatments, broadcast bait treatments and broadcast insecticide treatments. Individual mound treatments (e.g., liquid drenches or granules) are fairly fast acting, but only work to kill the mounds that are directly treated. They are most appropriate for eliminating individual mounds that present a hazard or as clean-up treatments. They are not effective as a management strategy for reducing the RIFA population in an area.</p> <p>Broadcast bait treatments are more effective in reducing the population in a given area. The speed at which the baits reduce mound numbers differs by products. Some are slower acting and no results will be noticed for three or four weeks. More recently introduced products act faster. Baits will most likely need to be applied in the spring and fall. Most baits are applied at a rate of 1 to 1.5 lbs per acre. The key to success with RIFA bait products is to broadcast the material when the ants are foraging since most of the products breakdown quickly in sunlight and water. Foraging activity is regulated by surface soil temperatures. RIFA forage when soil temperature reaches above 65°F. The best way to determine if ants are actively foraging is to place a small amount of test bait in the area to be treated. If RIFA hit the bait within 30 minutes then it is a good time to use the baits.</p> <p>Broadcast insecticide treatments are recommended for high use areas with zero tolerance for RIFA. Most of these applications are relatively expensive and therefore cost prohibitive in large areas. The advantage of the broadcast granular products is the high level of control that can be achieved. They are also relatively easy to use and can go out at any time of the year. Most of the products can give up to twelve months of control. To achieve good control, the product should be applied evenly and thoroughly to cover all areas to be treated.</p> <p>Combinations of IMT, bait, and/or granular broadcast treatments based upon the needs of the site are likely to produce better results than the use of a single chemical or strategy alone. The 'two-step' program is a combination of both broadcast bait and individual mound treatments, which is suitable for large and medium-sized area at a moderate cost. First step involves broadcasting slow-acting baits in the spring and/or fall, when the RIFA is actively foraging. Then individual RIFA mounds can be treated individually with a fast-acting contact insecticide in step two. This shortcoming of this treatment program is the relatively long time required before an overall reduction in the RIFA population and mounds will be observed.</p> <p>Check label of individual insecticide for information on site uses, application method, yearly application limit and buffer zone restrictions.</p>	
	bifenthrin ²	Orthene TT&O 97	Drench: 0.15 oz/gal/mound		
		Onyx Pro	Broadcast: 0.16 to 0.32 fl oz		
		Talstar EZ Golf, Talstar GC Gran	Broadcast: 2.3 to 4.6 lbs		
		Talstar GC Flowable, Talstar One	Broadcast: 0.5 to 1.0 fl oz		
	carbaryl	Talstar F	Drench: 1 teaspoon/gal/mound Broadcast: 1.0 fl oz		
		Sevin 10G	Drench: 1 teaspoon/gal/mound Broadcast: 1 to 1.9 oz/mound		
		Dursban 50W	Broadcast: 2 lbs/acre Sod Broadcast: 16 lbs/acre Drench: 0.05 lb/gal/mound		
	chlorpyrifos	Dursban PRO	Broadcast: 1.5 fl oz Drench: 0.5 fl oz/gal/mound		
		Tempo Ultra SC, Tempo Ultra GC	Broadcast: 0.27 fl oz		
	cyfluthrin	Tempo 20 WP, Tempo Ultra WP	Broadcast: 10 grams		
		Tempo 20 WP GC, Power Pak	1 packet/7,800 sq. ft.		
		Tempo 20 WSP	1 packet/5,000 sq. ft.		
	deltamethrin	DeltaGard GC, DeltaGard T&O	Drench: 1.5 fl oz/gal/mound		
		DeltaGard G, DeltaGard GC Gran	Broadcast: 2 to 3 lbs		
	fipronil	Chipco TopChoice	Broadcast: 2 lbs		
Chipco Choice		Broadcast: 4.6 oz			
imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	Broadcast: 2.9 to 5.7 lbs Dry: 0.5 cup/mound			
	Allectus SC, Allectus GC SC	Broadcast: 1.32 to 1.65 fl oz SC Drench: 1/3 fl oz/gal/mound GC Drench: 2/3 fl oz/gal/mound			
lambda-cyhalothrin	Demand EZ	Broadcast: 13.6 to 28 ml			
	Demand CS, Scimitar CS	Broadcast: 3.4 to 7 ml			
permethrin	Astro	Broadcast: 0.4 to 0.8 fl oz			
spinosad	Conserve SC	Drench: 0.1 fl oz/gal/mound			
thiamethoxam	Meridian 25WG	Drench: 1 to 3 oz/100 gal 2-3 gal//mound			
thiamethoxam + lambda-cyhalothrin	Tandem	Broadcast: 14-28 fl oz/acre			

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Ants, Red Imported Fire	Broadcast baits			<p>Many baits also control other nuisance ant, such as harvester ants and big-headed ants. Check label for specific instructions.</p> <p>Do not exceed 4 applications per year.</p> <p>Not for use on sod farms. Do not exceed 4 applications per year.</p> <p>Mix with other baits.</p>
	abamectin	Varsity Fire Ant Bait	Broadcast: 1 lb/acre Mound: 5 to 7 tbsp/mound	
	fenoxycarb	Award Fire Ant Bait	Broadcast: 1 to 1.5 lbs/acre Mound: 1 to 3 tbsp/mound	
	fipronil	Maxforce FC	Broadcast: 1.5 to 5 lbs/acre Mound: 2 to 5 tbsp/mound	
	hydramethylnon	Amdro Pro Fire Ant Bait SiegePro Fire Ant Bait Maxforce G	Broadcast: 1 to 1.5 lbs/acre Mound: 2 to 5 tbsp/mound Broadcast: 1 to 2 lbs/acre Mound: 2 to 5 tbsp/mound	
	indoxacarb	Advion Fire Ant Bait	Broadcast: 1.5 lb/acre Mound: 4 tablespoons/mound	
	methoprene	Extinguish	Broadcast: 1 to 1.5 lbs/acre Mound: 3 to 5 tbsp/mound	
	metaflumizone	Siesta	Mound: 1-2 oz.mound Broadcast: 1-1.5 lbs/acre	
	methoprene + hydramethylnon	Extinguish Plus	Broadcast: 1.5 lbs/acre Mound: 2 to 5 tbsp/mound	
	pyriproxyfen	Distance Fire Ant Bait	Broadcast: 1 to 1.5 lbs/acre Mound: 1 to 4 tbsp/mound	
spinosad	Justice Fire Ant Bait	Broadcast: 2.5 to 5 lbs/acre Mound: 4-6 tbsp/mound		
Billbugs (adults)	bifenthrin ²	Onyx, OnyxPro	0.07 to 0.16 fl oz, see labels	<p>Billbug adults and larvae feed on the roots and stems of various turfgrasses, with bermudagrass, zoysiagrass and seashore paspalum being the most susceptible. Symptoms are often misdiagnosed as drought stress or disease. Symptoms first appear as scattered dead stems and later enlarge to small patch turning from yellow to brown. Straw-color dead grasses are easily pulled out with the hollowed stem break away from the crown. Fine, sawdust-like frass can be seen at the base. The affected turf, which appears drought-stressed, does not recover with watering. Damage usually shows up in mid- to late-summer (worst in August) during extended drought period. Soil remains firm, not spongy underfoot as with white grub or mole cricket infestations.</p> <p>Adults can be forced from the grass with a detergent or captured with a pitfall trap. Adults can be found from March to November in SC; the peak activity occurs in May and September. Treat for grubs soon after adult peak activity. Detection and treatment for larvae are similar to white grubs. Treat when adults and/or larvae are found and damage is apparent. Most materials, particularly those against the grubs, should be watered-in with 2-inch immediately after application.</p> <p>Check label of individual insecticide for information on site uses, application method, yearly application limit and buffer zone restrictions.</p>
		Talstar EZ Golf, GC Gran, PL	1.15 to 2.3 lbs	
	carbaryl	Talstar GC Flowable, One, F	0.25 to 0.5 fl oz	
		Sevin 10G	1.4 to 1.9 lbs	
	chlorpyrifos	Dursban 50W	2 to 4 lbs/acre	
		Dursban PRO	1.5 fl oz	
	clothianidin + bifenthrin	Aloft GC SC, LC SC	11.65 to 23.3 fl oz, see label	
		Aloft GC G, LC G	80 to 160 lbs, see label	
	cyfluthrin	Tempo (various SC formulation)	see label	
		Tempo 20 WSP, Power Pak	1 packet/5,000 to 7,800 sq. ft.	
deltamethrin	DeltaGard GC, DeltaGard T&O	0.6 to 0.9 fl oz		
	DeltaGard GC Gran, DeltaGard G	2 to 3 lbs		
imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	1.7 to 2.9 lbs		
	Allectus SC	0.4 to 1.65 fl oz		

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Billbugs (adults)		Allectus GC SC	0.9 to 1.65 fl oz	Labeled for bluegrass billbug.
	lambda-cyhalothrin	Demand EZ	28 ml	
		Demand G	3 to 4 lbs	
	thiamethoxam + lambda-cyhalothrin	Demand CS, Scimitar GC and CS Tandem	7 ml 0.32 to 0.64 fl oz	
Billbugs (grubs)	carbaryl	Sevin 80 WSP	10 lb/acre	Begin preventive treatment against larvae soon after the adults become active in the spring (usually in late April and May; monitor with pitfall traps). Larger grubs can also be controlled curatively (although less effectively) in early summer (when the grubs are in the soil) with methods similar to the white grubs. 0.5G is not for use on sod farms. 1 packet = 1.6 oz 2SC is not for uses in residential lawns. Allow 7 days before treatment and harvest of sods when using Mach 2. Dylox is not for uses in sod farms, seed productions and research.
		Sevin SL	6 fl oz	
	chlorpyrifos	Dursban 50W	2 to 4 lbs/acre	
		chlorantraniliprole	Acelepryn	
			Acelepryn granule	
	clothianidin	Arena .25G	1.84 to 3.67 lbs	
		Arena .5G	1.0 to 1.8 lbs	
		Arena 50 WDG	6.4 to 12.8 oz	
	clothianidin + bifenthrin	Aloft GC SC	11.65 to 19 fl oz	
		Aloft GC G	80 to 132 lbs	
		Aloft LC SC	11.65 to 23.3 fl oz	
		Aloft LC G	80 to 160 lbs	
	dinotefuran	Zylam 20SG	1 oz	
	imidacloprid ²	Merit 0.5 G	1.4 to 1.8 lbs	
		Merit 2 F	0.46 to 0.6 fl oz	
		Merit 75 WP	3 to 4 teaspoons	
		Merit WSP	1 packet/8,250-11,000 sq. ft.	
	imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	1.7 to 2.9 lbs	
		Allectus SC	0.4 to 1.65 fl oz	
		Allectus GC SC	0.9 to 1.65 fl oz	
	halofenozide	Mach 2 2SC	2.9 fl oz	
		Mach 2 1.5G	3 lbs	
	thiamethoxam	Meridian 0.33G	1.42 to 1.88 lbs	
Meridian 25WG		0.3 to 0.39 oz		
thiamethoxam + lambda-cyhalothrin	Tandem	0.32-0.64 fl oz		
	trichlorfon	Dylox 6.2G	3 lbs	
		Dylox 80 T&O	3.75 fl oz/100 gal	
Caterpillars	acephate	Orthene TT&O 75, 97	see label	

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Caterpillars (Armyworms, Cutworms, Sod Webworms)	<i>Bacillus thuringiensis (Bt)</i>	Biobit, Crymax, Deliver, Dipel, Javelin, Lepinox, XenTari	0.5 to 3 lbs/acre, see label	
	bifenthrin ²	Onyx OnyxPro Talstar EZ Golf , GC Gran, PL Talstar GC Flowable, One, F	0.07 to 0.15 fl oz 0.05 to 0.08 fl oz 1.15 lbs 0.18 to 0.25 fl oz	Fall armyworm, cutworm and sod webworm are the major pest caterpillar species of turfgrass. All lawn caterpillars feed on grass blades.
	carbaryl	Sevin 10G Sevin 80 WSP Sevin SL	1.4 to 1.9 oz, see label 2.5 to 10 lb/acre, see label 1.5 to 6 fl oz, see label	True armyworms feed in groups, creating circular patches of bare grounds. Fall armyworms are less gregarious, so the damage often occurs as thinning of grasses in a broad area. Fall armyworm migrates from FL and southern GA each year. Damage begins to appear in June but at this time the caterpillars have become too big and difficult to control. Symptoms of care grounds often appear near building, trees, posts, poles and other erected structures as adult moths often lay eggs on these structures.
	chlorpyrifos	Dursban 50W Dursban PRO	2 lbs/acre 1.5 fl oz	Cutworms dig a burrow in the thatch or soil and eat the grass around the burrow at night, resulting in small patches of dead greens and sunken areas; thus reducing the smoothness and uniformity of putting surface. Frequent mowing and clipping removal help to reduce population but not enough to achieve control.
	chlorantraniliprole	Acelypryn Acelepryn granule	0.046 to 0.092 fl oz 1.15 to 2.3 lbs	
	clothianidin	Arena .25G Arena .5G Arena 50 WDG	1.84 to 3.67 lbs 1.0 to 1.8 lbs 12.8 oz	There are several generations of sod webworm in a year. The damage begins to appear in spring and becomes progressively severe with the season. Damage begins as general thinning, followed by small patches of brown, closely-cropped grass; later coalesces into large irregular patches with severe infestations. Adult sod webworm or lawn moths, which have characteristic snout-like projections in front of their heads, are active flyers over turfgrass in the evening.
	clothianidin + bifenthrin	Aloft GC SC, LC SC Aloft GC G Aloft LC G	11.65 to 23.3 fl oz; see label 80 to 132 lbs 80 to 160 lbs	The presence of birds feeding on the caterpillar should also be an indicator. All caterpillar species can be monitored with detergent flush.
	cyfluthrin	Tempo Ultra SC, Tempo Ultra GC Tempo 20 WP, Tempo Ultra WP Tempo 20 WP GC, Power Pak Tempo 20 WSP	0.135 to 0.27 fl oz 5 to 10 grams 1 packet/7,500-11,000 sq. ft. 1 packet/5,000-10,000 sq. ft.	Treated areas (with most contact insecticides) should not be irrigated within 24 hours of treatment so that the caterpillars will come in contact with the residues.
	deltamethrin	DeltaGard T&O, DeltaGard GC DeltaGard G, DeltaGard GC Gran	0.2 to 0.4 fl oz 2 to 3 lbs	Caterpillars are often attacked by natural enemies; therefore, conserve these natural enemies with less frequent sprays or compatible insecticides whenever possible. Use higher rates of Sevin for cutworm control.
	dinotefuran	Zylam 20SG	1 oz	Zylam is registered for use on cutworm and sod webworm.
	nametodes ²	Various products	See label	Only effective against small caterpillars.
	halofenozide	Mach 2 2SC Mach 2 1.5G	1.5 fl oz 1.55 lbs	Mach 2 for use in commercial sites only. Both formulations are more effective against younger caterpillars. They are preventive rather than curative in nature.
	imidacloprid ²	Merit 0.5 G Merit 2 F Merit 75 WP Merit 75 WSP	1.4 to 1.8 lbs 0.46 to 0.6 fl oz 3 to 4 teaspoons 1 packet/8,250-11,000 sq. ft.	Merit is for cutworm only. Must be applied against early stages. May only achieve suppression of the population. 1 packet = 1.6 oz.
	imidacloprid + bifenthrin	Allectus G, Allectus GC Gran Allectus GC SC	1.2 to 2.9 lbs 0.67 to 1.65 fl oz	

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments	
Caterpillars (Armyworms, Cutworms, and Sod Webworms)	indoxacarb	Allectus SC	0.4 to 1.65 fl oz	Not for use on sod farms.	
		Provaunt	0.046 to 0.092 oz		
	lambda-cyhalothrin	Demand EZ	13.6 to 28 ml		
		Demand G	2 to 3 lbs		
		Demand CS, Scimitar GC, CS	3.4 to 7 ml		
		permethrin	Astro		0.4 to 0.8 fl oz
		spinosad	Conserve SC		0.25 to 1.2 fl oz, see label
		thiamethoxam + lambda-cyhalothrin	Tandem		0.32-0.64 fl oz
trichlorfon	Dylox 80 T&O	2.5 to 3.75 fl oz/100 gal			
	Dylox 6.2G	2 lbs			
Chinch Bugs	acephate	Orthene TT&O 75 and 97	0.9 to 2.4 oz; see label	Chinch bugs are a major problem of St. Augustinegrass. Hot, dry weather, deep thatch, and high fertility favor chinch bug development. Infested turfgrass first appear yellow and stunted. As the infestation progresses, the grass wilt and dead, creating small dead patches which will expand as the infestation continues and the damage worsens. Sampling for chinch bugs is conducted using the floating method. Plugs of turf and soil can be placed in a 5-gallon bucket filled with clean water or an open ended cylinder (metal can or PIV pipe) driven into the ground and filled with clean water. Adults and nymphs will float to the surface within 10 minutes. Do not use soapy water. For best result sample along the edge of the damage. If chinch bugs are suspected and floating yields no results, visual examination of the stolens in the thatch layer may yield results. Treatments should be applied if 25 to 30 insects are found per square foot. Chinch bugs are often found in the thatch layer. Thus higher volume is critical in delivering the insecticides through the thatch layer and to successful insecticide treatments. See insecticide label for specifics. In general, use a minimum spray volume of 50 gallons/acre (1.2 gallons/1000 sq. ft.) . St. Augustinegrass varieties 'Floritam', 'Floralawn' and 'Captiva' are resistant to chinch bugs. Cultural controls include less N, using water insoluble (slow release) N, using a sharp mower blade, mowing at 3" in sun areas, 4" in shaded areas, and controlling thatch. Irrigate with ¾-inch when grass begins to wilt. Minimize the use of atrazine on St. Augustinegrass during summer. Monitor turf regularly. To preserve beneficial arthropods, limit treatment to the damaged area and 5 to 10 feet beyond. Recheck in 2-3 days. Spot treat again, if needed. Acelypryn provides suppression only. Zylam only achieves suppression. Merit only achieves suppression.	
	<i>Beauveria bassiana</i>	Botanigard	0.5 to 2 qts/100 gal		
	bifenthrin ²	Onyx, OnyxPro	See label		
		Talstar EZ Golf, GC Granular, PL	2.3 to 4.6 lbs		
	carbaryl	Talstar GC Flowable, One, F	0.25 to 0.5 fl oz		
		Sevin 10G	1.4 to 1.9 lbs		
		Sevin 80 WSP	7.5 to 10 lb/acre		
	chlorpyrifos	Sevin SL	4.5 to 6 fl oz		
		Dursban 50W	2 lbs/acre		
	chlorantraniliprole	Dursban PRO	1.5 fl oz		
		Acelypryn	0.184 to 0.46 fl oz		
	clothianidin	Acelepryn granule	1.15 to 2.3 lbs		
		Arena .25G and .5G	1.5 to 3.67 lbs, see label		
	clothianidin + bifenthrin	Arena 50 WDG	12.8 oz		
		Aloft GC SC	11.65 to 19 fl oz		
		Aloft GC G	80 to 132 lbs		
		Aloft LC SC	11.65 to 23.3 fl oz		
	cyfluthrin	Aloft LC G	80 to 160 lbs		
		Tempo (various formulations)	See label		
	cypermethrin	Demon Max	0.33 to 0.65 fl oz		
DeltaGard GC, DeltaGard T&O SC		0.6 to 0.9 fl oz			
deltamethrin	DeltaGard G, GC Gran, T&O Gran	2 to 3 lbs			
	dinotefuran	Zylam 20SG	1 oz		
imidacloprid ²	Merit 0.5 G	1.8 lbs			

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Chinch Bugs		Merit 2 F	0.6 fl oz	Meridian may only provide suppression.
		Merit 75 WP	4 teaspoons	
	imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	1.7 to 2.9 lbs	
		Allectus SC, Allectus GC SC	1.65 fl oz	
	lambda-cyhalothrin	Demand EZ	56 ml	
		Demand G	3 to 4 lbs	
		Demand CS, Scimitar GC, CS	14 ml	
permethrin	Astro	0.4 to 0.8 fl oz		
thiamethoxam	Meridian 0.33G	1.42 to 1.88 lbs		
	Meridian 25WG	0.3 to 0.39 oz		
thiamethoxam + lambda-cyhalothrin	Tandem	0.32 to 0.64 fl oz		
trichlorfon	Dylox 80 T&O	3.75 fl oz/100 gal		
Chiggers and Ticks	bifenthrin ²	OnyxPro	0.16 to 0.32 fl oz	Chiggers (red bugs) and ticks may be present in turfgrass areas. Although they do not damage turfgrass, they are however significant health risks. Insecticide treatments are often ineffective. Only DeltaGard, Demand, Demon, Dursban, Scimitar, Sevin and Tempo are labeled at higher rates for chigger control. Keep grass mowed short to discourage chiggers and ticks. Avoid contact by applying repellents and wearing protective clothing.
		Talstar EZ Golf, GC Gran, PL	2.3 to 4.6 lbs	
		Talstar GC Flowable, One, F	0.5 fl oz	
	carbaryl	Sevin 10G	1.4 to 1.9 lbs	
		Sevin 80 WSP	2.5 to 5 lb/acre	
		Sevin SL	1.5 to 3 fl oz	
	chlorpyrifos	Dursban 50W	2 lbs/acre	
		Dursban PRO	1.5 fl oz	
	cyfluthrin	Tempo Ultra SC, Tempo Ultra GC	0.135 to 0.27 fl oz	
		Tempo 20 WP, Tempo Ultra WP	5 to 10 grams	
	cyfluthrin	Tempo 20 WP GC, Power Pak	1 packet/7,500-11,000 sq. ft.	
		Tempo 20 WSP	1 packet/5,000-10,000 sq. ft.	
	deltamethrin	DeltaGard T&O, DeltaGard GC	0.4 to 0.6 fl oz	
		DeltaGard G, DeltaGard GC Gran	2 to 3 lbs	
	fipronil	Chipco TopChoice	2 lbs	
		Chipco Choice	4.6 oz	
	imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	2.9 lbs	
	Allectus GC SC, Allectus SC	1.32 to 1.65 fl oz		
lambda-cyhalothrin	Demand EZ	13.6 to 28 ml		
	Demand CS, Scimitar GC, CS	3.4 to 7 ml		
permethrin	Astro	0.4 to 0.8 fl oz		
			Broadcast application. 1 month control for ticks.	

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
	thiamethoxam + lambda-cyhalothrin	Tandem	0.24 fl oz	0.34 to 1.38 fl oz/1,000 sq ft for ticks.
Earthworms	No control is recommended.			Earthworms are considered beneficial organisms as they help aerate soil and decompose organic materials (such as thatch). However, when a large number of dirt and castings are pushed up to the putting greens, they will interfere with the play. At this time, no control is recommended. Earthworm castings may be managed with applications of saponin-containing products, such as the organic fertilizer Early Bird 3-0-1.
Fleas	acephate	Orthene TT&O 75	1.2 to 2.4 oz	Flea bites result in severe irritation and discomfort for human and pets. Pet owners can prevent infestation by treating the pets with a long-lasting insecticide or with flea collars. When curative treatment is needed, both infested area and pets need to be treated. Mowing the lawn before treatment may increase effectiveness. Use higher rates of bifenthrin for larval control, see label.
		Orthene TT&O 97	0.9 to 1.8 oz	
	bifenthrin	Onyx, OnyxPro	0.07 to 0.32 fl oz, see labels	
		Talstar EZ Golf , GC Gran, PL	2.3 to 4.6 lbs	
		Talstar GC Flowable, One, F	0.25 to 0.5 fl oz	
	carbaryl	Sevin 10G	1.4 to 1.9 lbs	
		Sevin 80 WSP	10 lb/acre	
		Sevin SL	6 fl oz	
	chlorpyrifos	Dursban 50W	2 lbs/acre	
		Dursban PRO	1.5 fl oz	
	cyfluthrin	Tempo Ultra SC, Tempo Ultra GC	0.27 fl oz	
		Tempo 20 WP, Tempo Ultra WP	10 grams	
		Tempo 20 WP GC, Power Pak	1 packet/7,800 sq. ft.	
		Tempo 20 WSP	1 packet/5,000 sq. ft.	
	cypermethrin	Demon Max, TC	0.33 to 0.65 fl oz	
	deltamethrin	DeltaGard T&O, DeltaGard GC	0.4 to 0.6 fl oz	
		DeltaGard G, DeltaGard GC Gran	2 to 3 lbs	
	fipronil	Chipco TopChoice	2 lbs	
	imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	2.9 lbs	
		Allectus SC, Allectus GC SC	0.4 to 1.65 fl oz, see label	
	lambda-cyhalothrin	Demand EZ	28 ml	
		Demand CS, Scimitar GC, CS	3.4 to 7 ml	
	permethrin	Astro	0.4 to 0.8 fl oz	
	spinosad	Conserve SC	1.2 fl oz	Target flea larvae.
	thiamethoxam + lambda-cyhalothrin	Tandem	0.34 to 1.38 fl oz	Perimeter treatment.
Grasshoppers	acephate	Orthene TT&O 75	0.5 oz	Grasshoppers are occasional pests of turfgrass. Usually infestations do not require treatment. During severe infestation, most contact insecticides are very effective.
		Orthene TT&O 97	0.4 oz	
	bifenthrin ²	Onyx	0.07 to 0.15 fl oz	

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Grasshoppers		OnyxPro	0.08 to 0.16 fl oz	Not for sod farms.
		Talstar GC Flowable, One, F	0.25 to 0.5 fl oz	
	carbaryl	Sevin 5 Bait	11 oz	
		Sevin 10G	1.4 to 1.9 oz	
		Sevin 80 WSP	2.5 to 5 lb/acre	
		Sevin SL	1.5 to 3 fl oz	
	clothianidin + bifenthrin	Aloft GC SC	11.65 to 19 fl oz	
		Aloft GC G	80 to 132 lbs	
		Aloft LC SC	11.65 to 23.3 fl oz	
		Aloft LC G	80 to 160 lbs	
	chlorpyrifos	Dursban 50W	2 lbs/acre	
		Dursban PRO	1.5 fl oz	
	cyfluthrin	Tempo Ultra SC, Tempo Ultra GC	0.135 to 0.27 fl oz	
		Tempo 20 WP, Tempo Ultra WP	5 to 10 grams	
		Tempo 20 WP GC, Power Pak	1 packet/7,500-11,000 sq. ft.	
	deltamethrin	DeltaGard T&O, DeltaGard GC	0.4 to 0.6 fl oz	
DeltaGard G, DeltaGard GC Gran		2 to 3 lbs		
imidacloprid + bifenthrin	Allectus SC	0.4 to 1.65 fl oz		
	Allectus GC SC	0.9 to 1.65 fl oz		
indoxacarb	Provaunt	0.275 oz		
lambda-cyhalothrin	Demand EZ	13.6 to 28 ml		
	Demand G	2 to 3 lbs		
	Demand CS, Scimitar GC, CS	3.4 to 7 ml		
thiamethoxam + lambda-cyhalothrin	Tandem	0.24 fl oz		
Greenbugs (Aphids)	acephate	Orthene TT&O 75	0.5 oz	
		Orthene TT&O 97	0.4 oz	
	bifenthrin ²	Onyx, Talstar	See label	
	carbaryl	Sevin	See label	
	chlorpyrifos	Dursban	See label	
	clothianidin	Arena .25G	See label	
	clothianidin + bifenthrin	Aloft	See label	
	cyfluthrin	Tempo	See label	

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Greenbugs (Aphids)	cypermethrin	Demon	See label	Currently, only Orthene is labeled specifically for greenbug control on golf courses and sod farms (not in landscape). Other contact and systemic insecticides are labeled for control of aphids on landscape ornamentals.
	imidacloprid + bifenthrin	Allectus	See label	
	imidacloprid ²	Merit	See label	
	lambda-cyhalothrin	Demand, Scimitar	See label	
	permethrin	Astro	See label	
	thiamethoxam thiamethoxam + lambda-cyhalothrin	Meridian Tandem	See label See label	
Leafhoppers	acephate	Orthene TT&O 75	1 oz	Leafhoppers can sometime become a significant pest of turfgrass. Leafhoppers overwinter as eggs or adults; with 1-5 generations per year. The adults and nymphs suck sap from grass blades, often causing silvery or whitish flecks or spots. Heavy infestation may cause mottled turf. Leafhopper infestations usually do not require treatment. When necessary, most contact and systemic insecticides are effective. Because of the movement of leafhoppers, repeated applications may be needed. Keep lawns and turf area well maintained. Healthy lawns can outgrow the damage.
		Orthene TT&O 97	0.75 oz	
	bifenthrin ²	Talstar EZ Golf , GC Gran, PL	1.15 to 2.3 lbs	
		Talstar GC Flowable, One, F	0.25 to 0.5 fl oz	
	carbaryl	Sevin 10G	1.4 to 1.9 lbs	
		Sevin 80 WSP	2.5 to 5 lb/acre	
		Sevin SL	1.5 to 3 fl oz	
	chlorpyrifos	Dursban 50W	2 lbs/acre	
		Dursban PRO	1.5 fl oz	
	clothianidin + bifenthrin	Aloft GC SC	11.65 to 19 fl oz	
		Aloft GC G	80 to 132 lbs	
		Aloft LC SC	11.65 to 23.3 fl oz	
		Aloft LC G	80 to 160 lbs	
	deltamethrin	DeltaGard T&O, DeltaGard GC	0.4 to 0.6 fl oz	
		DeltaGard G, T&O Gran, GC Gran	2 to 3 lbs	
imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	1.7 to 2.9 lbs		
	Allectus GC SC	0.9 to 1.65 fl oz		
	Allectus SC	0.4 to 1.65 fl oz		
lambda-cyhalothrin	Demand G	2 to 3 lbs		
Rhodesgrass Mealybugs	bifenthrin ²	Talstar EZ Golf , GC Gran, PL	1.15 to 2.3 lbs	Bermudagrass and St. Augustinegrass are most susceptible to attacks by the Rhodesgrass mealybug. Rhodesgrass mealybugs, similar to aphids and leafhoppers, feed by sucking the sap from leaf blades, stems and crowns. Damaged grass will first wilt, then turn from green to yellow to brown. The damage is especially serious during extended period of drought. When the grass is pulled up, the Rhodesgrass mealybugs are visible as white cottony masses attached to the nodes on grass stems. The mealybugs produced copious amount of honeydew, which is very attractive to ants and wasps. Normally Rhodesgrass mealybugs are not a problem for turfgrass, except in areas where the natural control by predators and parasites are disrupted. Conserve these natural
		Talstar GC Flowable, One, F	0.25 to 0.5 fl oz	
	carbaryl	Sevin	See label	
	chlorpyrifos	Dursban	See label	
	clothianidin	Arena	See label	
	clothianidin + bifenthrin cyfluthrin	Aloft Tempo	See label See label	

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Rhodegrass Mealybugs	deltamethrin	DeltaGard T&O, DeltaGard GC	0.4 to 0.6 fl oz	enemies by reducing the frequency of insecticide applications and using more compatible insecticides. Controlling fire ant may help reduce infestations.
	imidacloprid + bifenthrin	DeltaGard T&O Gran, GC Gran Allectus G, GC Gran Allectus SC, Allectus GC SC	2 to 3 lbs 1.7 to 2.9 lbs 0.9 to 1.65 fl oz	
	imidacloprid ²	Merit	See label	Currently, only Talstar, DeltaGard and Allectus are labeled specifically for mealybug control on turfgrass. Other contact and systemic insecticides are available for control of mealybugs in landscape ornamentals.
	lambda-cyhalothrin	Demand, Scimitar	See label	
	permethrin	Astro	See label	
	thiamethoxam thiamethoxam + lambda-cyhalothrin	Meridian Tandem	See label See label	
Millipedes Centipedes Pillbugs Sowbugs (Check cross reference table for specific chemicals)	acephate	Orthene TT&O 75	1.6 oz/gal	Millipedes, centipedes, pillbugs and sowbugs are common arthropods in turfgrass areas. They do not damage turfgrass. Centipedes may be important predators of other turfgrass pests. Control of these arthropods is not recommended. When necessary, perimeter treatment with contact insecticides can reduce the frequency of these arthropods invading buildings from the turfgrass areas. Higher rates for pillbug and sowbug control when using granules.
	bifenthrin ²	Talstar EZ Golf , GC Gran, PL	2.3 to 4.6 lbs	
	carbaryl	Talstar GC Flowable, One, F	0.25 to 0.5 fl oz	
		Sevin 10G	1.4 to 1.9 lbs	
		Sevin 80 WSP	2.5 to 5.0 lb/acre	
		Sevin SL	1.5 to 3 fl oz	
	chlorpyrifos	Dursban PRO	1.5 fl oz	
	clothianidin + bifenthrin	Aloft GC SC	11.65 to 19 fl oz	
		Aloft GC G	80 to 132 lbs	
		Aloft LC SC	11.65 to 23.3 fl oz	
		Aloft LC G	80 to 160 lbs	
	cyfluthrin	Tempo Ultra SC, Tempo Ultra GC	0.54 fl oz	
		Tempo 20 WP, Tempo Ultra WP	10 to 20 grams	
		Tempo 20 WP GC, Power Pak	1 packet/93-100 gal, see label	
	cypermethrin	Demon Max, TC	0.5 fl oz	
	deltamethrin	DeltaGard T&O, DeltaGard GC SC	0.4 to 0.6 fl oz	
		DeltaGard G, DeltaGard GC Gran, DeltaGard T&O Gran	2 to 3 lbs	
imidacloprid + bifenthrin	Allectus G, Allectus GC Gran	2.9-5.7 lbs		
	Allectus SC	0.4 to 1.65 fl oz		
	Allectus GC SC	0.9 to 1.65 fl oz		
lambda-cyhalothrin	Demand G	2 to 3 lbs		
	Demand EZ	13.6 to 28 ml		
	Demand CS	3.4 to 7 ml		
	Scimitar GC, CS	3.4 to 7 ml		

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments		
	permethrin	Astro	0.4 to 0.8 fl oz			
Mites	bifenthrin ²	Onyx OnyxPro Talstar GC Flowable, One, F	0.07 to 0.15 fl oz 0.08 to 0.16 fl oz 0.25 to 0.5 fl oz	<p>Clover mites are a problem in spring and fall. Feeding damage is a silvery discoloration of grass and often near the house foundation. Bermudagrass mites (an eriophyid mite) are a problem on common bermudagrass during hot, dry weather. Feeding causes yellowing and distortion of the grass, often results in a tufted or twisted 'rosette' or 'witch-broom' growth. Another common eriophyid mite pest of warm-season turfgrass is the zoysiagrass mite which causes rosette growth on infested zoysiagrass. Newer, hybrid bermudagrasses and zoysiagrass are resistant to their respective mite pests.</p> <p>Management of bermudagrass mite is especially problematic because no registered products are completely effective against this species. Diazinon has been shown in a study conducted in the 1980s to be the most effective active ingredient. Studies are currently underway to determine the best timing and materials to use for the management of bermudagrass mite.</p> <p>A wetting agent in the spray mixture improves control. Reapply in 10-14 days. Cultural controls include collecting and removing clippings. Reduce mowing height as close as practical if mites are a problem.</p> <p>Do not use Kelthane and Dicofof on residential lawns.</p>		
Mites (Clover, Bermudagrass, and other eriophyid mites)	chlorpyrifos	Dursban 50W Dursban PRO	2 lbs/acre 1.5 fl oz			
	cyfluthrin	Tempo 20 WP GC, Power Pak	1 packet/93-100 gal, see label			
	deltamethrin	DeltaGard GC, DeltaGard T&O DeltaGard GC Gran, DeltaGard G, DeltaGard T&O G	0.6 to 0.9 fl oz 2 to 3 lbs			
	dicofof	Kelthane 50 WSP Dicofof 4E	0.5 to 1 lb/acre 2/3 to 1 pints/acre			
	imidacloprid + bifenthrin	Allectus SC Allectus GC SC	0.4 to 1.65 fl oz 0.9 to 1.65 fl oz			
	lambda-cyhalothrin	Demand G Demand EZ Demand CS Scimitar GC, CS	2 to 3 lbs 13.6 to 28 ml 3.4 to 7 ml 3.4 to 7 ml			
	Mole Crickets	acephate	Orthene TT&O 75 Orthene TT&O 97		1.0 to 1.9 oz 0.8 to 1.4 oz	<p>Mole cricket adults are present during later winter and early spring. Mating flights occur from April through June. Egg hatch occurs from mid-June through July. The tawny mole cricket is a much more serious problem than the southern mole cricket.</p> <p>Tunneling is the most obvious sign of mole cricket infestation. To detect mole crickets, use a detergent flush consisting of 1 to 2 fl oz liquid detergent per gallon of water. One gallon will flush a 4 sq. ft. area. Treat when mole crickets and damage are present.</p> <p>Treatment in the early spring is probably beneficial because this will reduce the number of adult mole crickets laying eggs. Although small nymphs cause little noticeable damage, their treatment in late June and July is highly recommended. Sprays and granules should be applied during mid to late June. Application of baits and Orthene should be made when damage first appears (early- to mid-July). Insecticides can be applied later in the year (Aug.-Oct.). Soil should be moist at time of treatment. If soil is not moist, it is important to irrigate before applying sprays, granules and baits. After treatment, irrigate sprays or granulars into soil with 2 inch of water, except Orthene and baits. A surfactant may increase the efficacy of Orthene. Apply all pesticides as late in the day as possible. Do not irrigate after application of baits for 2-3 days if possible. Use a higher rate for large nymphs and adult mole cricket control.</p> <p>Cultural controls include not mowing turf shorter than recommended heights. Use a sharp mower blade. Maintain proper fertility and pH levels, as well as irrigation practices.</p>
		bifenthrin ²	Onyx OnyxPro Talstar EZ Golf , GC Gran, PL Talstar GC Flowable, One, F		0.07 to 0.15 fl oz 0.16 to 0.32 fl oz 2.3 to 4.6 lbs 0.25 to 1.0 fl oz	
		carbaryl	Sevin 5 Bait		20 to 48 lbs/acre	
	chlorpyrifos	Dursban 50W	4-6 lbs/acre			
	clothianidin	Arena .25G Arena .5G Arena 50 WDG	1.84 to 3.67 lbs 1.5 to 1.8 lbs 12.8 oz			
	clothianidin + bifenthrin	Aloft GC SC, LC SC Aloft GC G, LC G	11.65 to 23.3 fl oz, see label 80 to 160 lbs, see label			
	cyfluthrin	Tempo (various formulations)	See label			
	deltamethrin	DeltaGard GC 5 SC DeltaGard GC Gran, DeltaGard G	0.6 to 0.9 fl oz 2 to 3 lbs			

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments	
Mole Crickets	dinotefuran	Zylam 20SG	1 oz	Arena provides only suppression.	
	nematodes ²	Nematac S, various	See label	Slit application when using Chipco Choice.	
	fipronil	Chipco Choice 0.1 G TopChoice	4.6 to 9.4 oz 2 lbs		
	imidacloprid ²	Merit 0.5 G	1.8 lbs	Advion Mole Cricket Bait is not for use on sod farms and seed productions. See supplemental label.	
		Merit 2 F	0.6 fl oz		
		Merit 75 WP	4 teaspoons		
	imidacloprid + bifenthrin	Merit 75 WSP	1 packet/8,250 sq. ft.		2.9 to 5.7 lbs
		Allectus G, Allectus GC	1.32 to 3.3 fl oz		
	indoxacarb	Allectus SC, Allectus GC SC	1.15 to 4.6 lbs/acre		0.275 oz
		Advion Mole Cricket Bait	3 to 4 lbs		
lambda-cyhalothrin	Provanut	28 to 56 ml	7 to 14 ml		
	Demand G	0.4 to 0.8 fl oz			
	Demand EZ	1.42 to 1.88 lbs			
permethrin	Demand CS, Scimitar GC or CS	0.3 to 0.39 oz	0.32 to 0.64 fl oz		
	Astro	3.75 fl oz/100 gal.			
thiamethoxam	Meridian 0.33G	3 lbs	Meridian provides only suppression.		
	Meridian 25WG				
thiamethoxam + lambda-cyhalothrin	Tandem				
	trichlorfon	Dylox 80 T&O Dylox 6.2G			
Snails and Slugs	iron phosphate	Sluggo	1 lb	Apply late in the evening, especially after rain or irrigation. Reapply when needed. Water infested area thoroughly before application. Do not re-water for 48 hours.	
	iron phosphate + spinosad	Sluggo Plus	0.5 lb		
	mesurol	Mesurol 2% Bait	1 lb		
	metaldehyde	Deadline	6.4 oz		
Spittlebugs	acephate	Orthene TT&O 75	1.0 to 1.9 oz	Spittlebugs are increasingly damaging, particularly to centipedegrass. The most common pest species of turfgrass in South Carolina is the two-lined spittlebugs. Adult two-lined spittle bugs have a reddish black with 2 orange or red lines across the wings and a bright red abdomen. Nymphs are found at the base of the grass plant. The nymphs are enclosed individually in white foamy spittle masses. Feeding causes yellowing of the grass. High mowing height and thatch buildup aggravate the problem. There are typically two generations in SC: the first peak of adults occurs in June-July and the second peak in August-September. Overwinter as eggs.	
		Orthene TT&O 97	0.9 to 1.8 oz		
	bifenthrin ²	Onyx	0.07 to 0.15 fl oz		
		Sevin 10G	1.4 to 1.9 lbs		
	carbaryl	Sevin 80 WSP	2.5 to 5.0 lb/acre		
		Sevin SL	1.5 to 3 fl oz		

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
Spittlebugs	clothianidin	Arena .25G	1.84 to 3.67 lbs	<p>Treat when nymphs are present and damage appears. Use a minimum of 50 gallons of water per acre (1.2 gallons/1000 sq. ft.). Mow and irrigate before treatment. Treat also the ornamentals, particularly hollies, for adults.</p> <p>Dethatching , withholding irrigation when nymphs are hatching in May and July, and avoiding planting Japanese hollies near centipede grass can help reduce infestations.</p>
		Arena .5G	1.0 to 1.8 lbs	
	clothianidin + bifenthrin	Aloft GC SC	11.65 to 19 fl oz	
		Aloft GC G, LC G	80 to 160 lbs	
		Aloft LC SC	11.65 to 23.3 fl oz	
cyfluthrin	Tempo 20 WP GC, Power Pak	1 packet/93-100 gal, see label		
deltamethrin	DeltaGard G, GC Gran, T&O Gran	2 to 3 lbs		
lambda-cyhalothrin	Demand G	2 to 3 lbs		
Wasps, Bees (Burrowing or Digging)	acephate	Orthene TT&O 75	1.6 oz/gal	Mounds and nests made by soil burrowing wasps and bees can interfere with play and the appearance of the lawns and fields. Treatment should be done in the evening when foraging adults have returned to the nest. Thoroughly spray the entrance to the nest.
		Orthene TT&O 97	1.2 oz/gal	
	bifenthrin ²	Talstar One	0.25 to 0.5 fl oz	Burrowing and digging wasps and bees often dig in sandy areas. Improve the grass coverage or re-sodding the thinned areas will help to dissuade the wasps and bees from building a nest.
		Dursban 50W	0.25 to 4 lbs/50 gal, see label	
	chlorpyrifos	Dursban PRO	0.17 to 2.7 fl oz/gal, see label	Treatments of landscape ornamentals that are frequented by the honeybees and the native bees, with the hope of killing or dissuading visits by the bees, are strongly discouraged.
		Tempo Ultra SC, Tempo Ultra GC	0.54 fl oz/gal	
	cyfluthrin	Tempo 20 WP, Tempo Ultra WP	20 grams/gal	Lambda-cyhalothrin is registered for outdoor ornamental control of wasps.
		Demon Max, TC	1 fl oz/gal	
	deltamethrin	DeltaGard G, DeltaGard GC Gran	2 to 3 lbs	
	lambda-cyhalothrin	Demand	See label	
permethrin	Scimitar	See label		
	Astro	0.4 to 0.8 fl oz		
White Grubs	carbaryl	Sevin 10G	1.9 lbs	<p>May and June beetle, green June beetle, masked chafer, European chafer, black turfgrass ateniensis, oriental beetle and Japanese beetle larvae are all grouped under white grubs. Identification of the species is based on the raster (area of spines, hair and bare spaces on the underside of the last abdominal segment).</p> <p>White grubs live in the soil and feed on the grass roots. Infested turf often appears wilt and turns yellow and then brown. Heavily damaged turf feels spongy underfoot. Damaged sod easily fall apart when cut or lifted. Large green June beetle grubs also burrow to the surface at night and cause small dirt mounds on the turf.</p> <p>To detect grubs, remove 1 sq. ft. of sod and carefully inspect the root zone. Pay particular attentions to areas with high feeding and searching activity of black parasitic wasps and other animals. In most cases, if more than 7 grubs are found per sq. ft., treatment is needed. Apply preventive treatment (using neonicotinoids and chlorantraniliprole) in April – June or curative treatment against small grubs in July-August. Most materials should be watered-in immediately (at least 2 inches) after</p>
		Sevin 80 WSP	10 lb/acre	
		Sevin SL	6 fl oz	
	chlorpyrifos	Dursban 50W	2 to 8 lbs/acre	
		Dursban PRO	1.5 fl oz	
	chlorantraniliprole	Acelepryn	0.184 to 0.367 fl oz	
		Acelepryn granule	1.15 to 2.3 lbs	
	clothianidin	Arena .25G	1.84 to 3.67 lbs	
		Arena .5G	1.0 to 1.8 lbs	
		Arena 50 WDG	6.4 to 12.8 oz	
clothianidin + bifenthrin	Aloft GC SC	11.65 to 19 fl oz		

INSECT PEST CONTROL

Pest	Pesticide Common Name	Pesticide Trade Name and Formulation ¹	Rate / 1,000 sq. ft.	Pest Biology, Symptoms, Cultural Practices, and Comments
White Grubs	dinotefuran halofenozide	Aloft GC G	80 to 132 lbs	application. See label directions. Dursban Pro is only for green June beetle grub control.
		Aloft LC SC	11.65 to 23.3 fl oz	
		Aloft LC G	80 to 160 lbs	
		Zylam 20SG	1 oz	
		Mach 2 2SC	2.9 fl oz	
		Mach 2 1.5G	3 lbs	
	imidacloprid ²	Merit 0.5 G	1.4 to 1.8 lbs	Demand and Scimitar only provide suppression.
		Merit 2 F	0.4 to 0.6 fl oz	
		Merit 75 WP	3 to 4 teaspoons	
		Merit 75 WSP	1 packet/11,000 sq. ft.	
	imidacloprid + bifenthrin	Allectus G, Allectus GC	2.3 to 2.9 lbs	
		Allectus SC, Allectus GC SC	1.32 to 1.65 fl oz	
	lambda-cyhalothrin	Demand SC, Scimitar SC	7 ml	
		Demand EZ	28 ml	
		Demand G	3 to 4 lbs	
thiamethoxam	Meridian 0.33G	1.42 to 1.88 lbs		
	Meridian 25WG	0.3 to 0.39 oz		
thiamethoxam + lambda-cyhalothrin trichlorfon	Tandem	0.32 to 0.64 fl oz		
	Dylox 6.2G	3 lbs		
	Dylox 80 T&O	3.75 fl oz/100 gal		

¹Always check to be sure the formulation that you purchase is labeled for the site and pest you intend to use it for. No endorsement of products is intended, nor is criticism of unnamed products implied. **Read container label carefully for, use directions, application techniques, irrigation requirements, worker protection information, and precautions. Be sure** the formulation of pesticide you buy and use is labeled for use on turfgrass.

²Several trades names available. Check label for active ingredients, formulations and instructions.